

Comparing the Effectiveness of Emotional Disclosure Through Drawing and Family-Based Floortime Play Therapy on Emotion Regulation Difficulties and Internalizing and Externalizing Symptoms in Children with Anxiety Symptoms

Amir Dariush. Tavasoli¹, Samaneh Sadat. Jafar Tabatabaei^{2*}, Fatemeh. Shahabizadeh², Maryam. Nasri²

1 PhD Student, Department of Psychology, Bi.C., Islamic Azad University, Birjand, Iran.

2 Assistant Professor, Department of Psychology, Bi.C., Islamic Azad University, Birjand, Iran.

*Correspondence: Tabatabaee88@yahoo.com

Article type:
Original Research

Article history:
Received 14 July 2025
Revised 02 October 2025
Accepted 07 October 2025
Published online 01 January 2026

ABSTRACT

This study aimed to compare the effectiveness of emotional disclosure through drawing and family-based Floortime play therapy in improving emotion regulation and reducing internalizing and externalizing symptoms in children with anxiety symptoms. This quasi-experimental study used a pretest–posttest design with a control group and follow-up. The statistical population comprised children aged 8–10 years in Qaen, South Khorasan Province, in 2024–2025. Forty-five children with anxiety symptoms, identified by scoring at least one standard deviation above the mean on the Spence Children's Anxiety Scale–Parent version (SCAS-P), were selected through purposive sampling and randomly assigned to three groups: emotional disclosure through drawing, family-based Floortime play therapy, and control (15 participants each). Both interventions were delivered in eight 45-minute sessions held twice weekly. Data were collected using the Difficulties in Emotion Regulation Scale for Children (DERS-C) and the Child Behavior Checklist (CBCL) and analyzed with analysis of covariance (ANCOVA) and Bonferroni post hoc tests using SPSS 26. ANCOVA results indicated a significant main effect of group on emotion regulation difficulties ($F = 14.926$, $p = .001$, $\eta^2 = .421$), internalizing symptoms ($F = 30.363$, $p < .001$, $\eta^2 = .603$), and externalizing symptoms ($F = 36.531$, $p < .001$, $\eta^2 = .646$). Bonferroni comparisons showed both interventions were significantly more effective than the control group at posttest and follow-up ($p < .001$). No significant difference was observed between emotional disclosure and Floortime groups ($p > .05$). Both emotional disclosure through drawing and family-based Floortime play therapy effectively improved emotion regulation and reduced anxiety-related internalizing and externalizing symptoms in children, with sustained effects at follow-up, suggesting these methods as accessible and family-friendly therapeutic options.

Keywords: Emotional disclosure, Drawing therapy, Floortime play therapy, Emotion regulation, Internalizing symptoms, Externalizing symptoms, Child anxiety

How to cite this article:

Tavasoli, A.D., Jafar Tabatabaei, S., Shahabizadeh, F., & Nasri, M. (2026). Comparing the Effectiveness of Emotional Disclosure Through Drawing and Family-Based Floortime Play Therapy on Emotion Regulation Difficulties and Internalizing and Externalizing Symptoms in Children with Anxiety Symptoms. *Mental Health and Lifestyle Journal*, 4(1), 1-13. <https://doi.org/10.61838/mhlj.212>

Introduction

Childhood is a sensitive period for the development of emotional and behavioral competencies that underpin mental health and adaptive functioning. One of the most important challenges faced by children, especially those who present with anxiety symptoms, is difficulty in regulating emotions. Emotion regulation difficulties are consistently associated with increased vulnerability to a wide range of psychopathological outcomes, including anxiety, depression, and externalizing behavioral problems (1). In particular, children with impaired ability to recognize, express, and modulate emotions may struggle to form stable relationships, engage effectively in academic settings, and cope with environmental stressors (2, 3). These regulatory deficits can lead to the manifestation of internalizing problems, such as withdrawal and somatic complaints, and externalizing difficulties, such as aggression and rule-breaking behavior (4). As such, identifying effective interventions to enhance emotional regulation and reduce behavioral symptoms is a key priority in contemporary child psychology and education (5).

Anxiety disorders and subclinical anxiety symptoms are highly prevalent in childhood and early adolescence and frequently co-occur with emotion dysregulation (6). Anxiety is not merely a transient developmental challenge but can predict long-term maladjustment if left untreated. Children experiencing persistent worry, avoidance, and somatic tension are at greater risk of impaired social functioning and academic underachievement (3, 7). Moreover, family relational patterns, including parental stress and maladaptive emotional expression, may exacerbate these vulnerabilities (8). Therefore, modern therapeutic approaches emphasize family involvement and experiential methods that engage children in a developmentally appropriate, emotionally safe way (9, 10).

One promising approach for addressing emotional and behavioral dysregulation is Floortime play therapy, also known as the Developmental, Individual-differences, Relationship-based (DIR) model. Originally designed for children with autism spectrum disorders, Floortime focuses on strengthening parent–child emotional connections, scaffolding self-regulation, and promoting flexible social engagement through play (11). Research shows that adapting Floortime to various developmental and emotional challenges beyond autism yields positive results. For instance, Floortime interventions have improved social skills and communication in neurodiverse populations (9, 12), enhanced executive functions and cognitive emotion regulation in children with ADHD (13), and increased social adaptability in children with emotional and behavioral issues (10, 14). Its family-based format strengthens caregivers' ability to interpret and respond to the child's affective signals, helping reduce parenting stress and support a secure emotional environment (15).

Another intervention with growing empirical support is emotional disclosure through drawing. Expressive writing has long been recognized as a method for processing traumatic and distressing experiences (16), but children often lack the verbal sophistication needed for such narrative work. Art-based emotional disclosure allows children to externalize painful or confusing feelings in a symbolic and developmentally appropriate way (17). By using drawings, children can represent complex inner states safely and tangibly, facilitating both catharsis and cognitive reorganization of distressing memories (18). Evidence indicates that drawing-based disclosure can reduce aggression and anger (18), alleviate stress in parents of children with developmental challenges (15), and contribute to improvements in emotional processing and resilience (19).

Recent adaptations integrate digital art to further enhance engagement and self-expression among children (17).

The combination of art-based disclosure and play-based relational interventions reflects a broader movement toward experiential, child-centered therapies. These modalities are particularly well-suited for younger children, who may have difficulty with abstract discussion of emotions and instead benefit from tangible, action-oriented strategies (9, 20). For example, integrating drawing and group play has been shown to reduce both internalizing and externalizing problems in children with separation anxiety disorder (20). Likewise, play therapies that actively engage parents not only address child-level symptoms but also target family-level factors such as emotional climate and parental responsiveness (9, 15). Such family-based interventions can indirectly reduce maladaptive parentification, perfectionism, and suppression of emotion in both children and their caregivers (7).

From a developmental psychopathology perspective, internalizing and externalizing symptoms often share underlying vulnerabilities in emotion regulation systems (1, 4). Poor modulation of affect can lead to internal distress, such as anxiety and depression, or to outward behavioral difficulties, depending on environmental contingencies and individual temperament (2, 3). Consequently, treatments that strengthen emotional awareness, acceptance, and self-regulatory strategies may have broad transdiagnostic benefits (5). Transdiagnostic approaches that address common affective mechanisms are increasingly recognized as cost-effective and clinically powerful (4). Floortime's emphasis on emotional signaling and relational repair aligns well with this transdiagnostic model, as does expressive art therapy, which fosters both affect labeling and nonverbal processing of distress (17).

Family dynamics also play a critical role in shaping children's emotional competencies. High levels of expressed emotion, parental criticism, and inconsistent support can reinforce maladaptive schemas and impede healthy affect regulation (8). Conversely, interventions that train caregivers to attune to children's cues and respond contingently can transform these relational patterns (9, 10). Family-centered Floortime not only strengthens dyadic attunement but also reduces parenting stress, thereby creating a more stable emotional environment (15). When combined with child-centered emotional disclosure through drawing, which allows children to safely process and reorganize painful experiences, these interventions address both systemic and individual pathways to improved mental health (20).

The relevance of these therapeutic methods extends beyond clinical populations, as emotion regulation competence is a key predictor of academic functioning, social integration, and resilience under stress (3, 7). Effective early interventions can prevent escalation of subclinical anxiety into more entrenched disorders and reduce long-term service needs (6). Furthermore, creative therapies may have cultural advantages, providing a non-stigmatizing entry point for families and children reluctant to engage in traditional talk therapy (19). The adaptability of art-based and play-based methods across cultural contexts, including Iran, has been supported by local research demonstrating positive outcomes for emotional regulation and behavioral adjustment (9, 20).

Despite their promise, comparative studies examining the relative effectiveness of emotional disclosure through drawing and family-based Floortime in children with anxiety symptoms remain scarce. Much of the existing evidence evaluates each intervention in isolation or in populations with autism spectrum disorder or behavioral difficulties, but less is known about their direct impact on anxiety-driven internalizing and

externalizing symptoms. Moreover, the potential maintenance of therapeutic gains over time—a critical aspect of clinical effectiveness—has been underexplored (5).

Given the need for accessible, developmentally sensitive, and family-inclusive interventions, this study aimed to compare the effectiveness of emotional disclosure through drawing and family-based Floortime play therapy in reducing emotion regulation difficulties and alleviating internalizing and externalizing symptoms in children with anxiety symptoms. By directly contrasting these two promising approaches, the research seeks to inform clinical decision-making and provide evidence for targeted, culturally relevant therapeutic strategies to support children's emotional well-being and adaptive functioning.

Methods and Materials

Study Design and Participants

This study employed a quasi-experimental design with a pretest–posttest control group and a follow-up period. The statistical population included all children aged 8 to 10 years residing in Qaen, South Khorasan Province, in 2024–2025. Children with anxiety symptoms were identified as those who met the study inclusion criteria and scored more than one standard deviation above the mean on the Spence Children's Anxiety Scale–Parent version (SCAS-P), which helps accurately identify children with elevated social anxiety and related internalizing problems (mean = 36.76, SD = 5.63).

Initially, 200 children were screened using the SCAS-P. Based on Cohen's table for comparing three groups with a medium effect size, a minimum statistical power of 0.80, and a significance level of 0.05, a sample size of 15 participants per group was determined. Therefore, a total of 45 participants were selected through purposive sampling after verifying eligibility criteria and were randomly assigned to three groups: emotional disclosure through drawing, family-based Floortime play therapy, and a control group.

Inclusion criteria were the absence of chronic physical illnesses, confirmed elevated anxiety symptoms on the SCAS-P, and the willingness of families to actively participate in family-based Floortime sessions. Exclusion criteria included absence from more than two intervention sessions and failure to provide signed informed consent. Parents provided written informed consent after receiving full explanations of the study aims and procedures.

Both intervention groups underwent eight 45-minute sessions. The emotional disclosure group engaged in guided drawing activities designed to encourage expression of feelings and experiences. The family-based Floortime group participated in structured play sessions involving the child, parent(s), and therapist, aiming to enhance emotional connection, self-regulation, and communication. The control group was placed on a waitlist and, after the study follow-up, was offered two free educational sessions to ensure ethical fairness.

Data Collection

The Difficulties in Emotion Regulation Scale for Children (DERS-C), originally developed by Gratz and Roemer (2004) and adapted for children, was used to measure emotion regulation difficulties. The revised 36-item version assesses six dimensions: nonacceptance of emotional responses, difficulties engaging in goal-directed behavior, impulse control difficulties, lack of emotional awareness, limited access to emotion regulation strategies, and lack of emotional clarity. Items are scored on a Likert scale, with higher scores reflecting greater difficulties in emotion regulation. The DERS-C demonstrates strong internal consistency,

with Cronbach's alpha reported at 0.93 for the full scale and above 0.80 for each subscale. In Iranian validation studies, Cronbach's alpha coefficients ranged from 0.82 to 0.89, and test–retest reliability ranged between 0.79 and 0.91 over a one-week interval. In the present study, internal consistency was 0.89.

Behavioral and emotional problems were assessed using the Child Behavior Checklist for ages 6–18 (CBCL) developed by Achenbach (2001). The parent-report form evaluates competencies, adaptive functioning, and a broad range of emotional and behavioral difficulties. It includes 115 items rated on a 3-point Likert scale from 0 (not true) to 2 (very true/often true). The CBCL measures eight syndrome scales: anxious/depressed, withdrawn/depressed, somatic complaints, social problems, thought problems, attention problems, rule-breaking behavior, and aggressive behavior. It also provides three broadband scores: internalizing problems, externalizing problems, and total problems. T scores below 60 are considered nonclinical, scores from 60 to 63 (or up to 65 for syndrome scales) are borderline clinical, and higher scores indicate a clinical range. The CBCL has demonstrated excellent reliability and validity across studies, with Cronbach's alpha up to 0.97 and test–retest reliability of 0.94. It has been standardized and validated for use in Iran, with strong internal consistency and construct validity confirmed in local samples.

Interventions

The family-based Floortime play therapy program was implemented in eight 45-minute sessions held twice weekly. The intervention followed the structured protocol developed by Zarrabi Moghadam and Yazdi (2019) and was delivered by the researcher. The sessions began with building rapport, introducing the therapy space, and familiarizing mothers and children with the principles and rules of Floortime play therapy. Mothers were guided to observe their children's unique sensory and emotional responses, understand individual differences, and engage in responsive and emotionally attuned interactions. As the program progressed, sessions focused on strengthening mother–child communication through guided play, encouraging children to lead activities while therapists and mothers scaffolded self-expression, impulse control, and goal-directed behavior. Activities included interactive games tailored to sensory processing, memory, and problem-solving, such as hiding and guessing objects, creating stories, drawing without certain letters, and symbolic play to improve emotional connection and cognitive flexibility. Mothers were taught to recognize and respond to their children's emotional cues, close communication loops, and reinforce adaptive self-talk. By the final sessions, the focus was on integrating acquired skills, enhancing mutual understanding, and helping children link ideas and emotions, while parents were coached to maintain therapeutic play at home. Each child received a commemorative gift at the conclusion of the program, and posttest assessments were conducted immediately after the last session.

The emotional disclosure through drawing program consisted of eight 45-minute sessions conducted twice weekly, based on the intervention framework of Ashour and colleagues (2023) and delivered by the researcher. The sessions started with establishing a safe therapeutic environment, explaining group rules in child-friendly language, and encouraging participants to express unpleasant or stressful memories through drawings that captured their deepest emotions and thoughts. In subsequent sessions, children expanded and reinterpreted their previous drawings to deepen their emotional expression without needing to recount specific events verbally. As therapy advanced, the focus shifted to enhancing social understanding and relationship building by assigning symbolic colors to interpersonal experiences and drawing facial

expressions representing varied emotions such as happiness, sadness, anger, and calmness. Children were guided to reflect on their own and others' emotional states, examine the impact of hasty judgments on relationships, and explore healthier ways of responding. Later sessions targeted anger management and self-control by drawing aggressive and calm animals, comparing their appeal, and discussing personal choices. In the final meetings, children transformed negative and positive emotion words into images of people, animals, or objects, gradually recognizing their own emotional strengths and areas for growth. The closing session involved sharing these drawings with the group, fostering self-acceptance, and reinforcing the therapeutic gains before posttest assessment.

Data analysis

Data were analyzed in two stages: descriptive and inferential. Descriptive statistics, including means and standard deviations, were calculated for demographic variables (age, gender) and outcome measures at pretest, posttest, and follow-up stages. Inferential analyses were performed using SPSS version 26. To examine the effects of the interventions, analysis of covariance (ANCOVA) was employed, controlling for pretest scores to compare posttest outcomes across groups. Bonferroni post hoc tests were used to identify pairwise differences between the emotional disclosure, Floortime play therapy, and control groups. Prior to conducting ANCOVA, assumptions such as homogeneity of regression slopes, normality, and homogeneity of variances (Levene's test) were tested and confirmed. The statistical significance level was set at $p < .05$ for all analyses.

Findings and Results

The final study sample consisted of 45 children aged 8 to 10 years, distributed equally across the three groups of emotional disclosure through drawing, family-based Floortime play therapy, and control. In the emotional disclosure group, 2 children (4.4%) were 8 years old, 7 children (15.5%) were 9 years old, and 6 children (13.3%) were 10 years old. In the Floortime group, 1 child (2.2%) was 8 years old, 6 children (13.3%) were 9 years old, and 8 children (17.7%) were 10 years old. In the control group, 2 children (4.4%) were 8 years old, 6 children (13.3%) were 9 years old, and 7 children (15.5%) were 10 years old. Overall, the sample showed a relatively balanced age distribution, with the largest proportion of participants being 9 and 10 years old across all groups, ensuring comparability among intervention and control conditions.

Table 1. Descriptive indices of emotion regulation difficulties and internalizing and externalizing symptoms across groups

Variable	Group	Stage	Mean	SD	Mean	SD	Mean	SD
Emotion Regulation Difficulties	Emotional Disclosure Drawing	Pretest	116.47	2.99	Posttest	70.13	1.60	Follow-up
	Floortime Play Therapy	Pretest	115.13	3.16	Posttest	73.06	2.05	Follow-up
	Control	Pretest	110.80	3.08	Posttest	90.06	1.79	Follow-up
Internalizing Symptoms	Emotional Disclosure Drawing	Pretest	88.06	12.40	Posttest	54.00	4.94	Follow-up
	Floortime Play Therapy	Pretest	83.00	11.03	Posttest	52.12	4.68	Follow-up
	Control	Pretest	88.33	13.98	Posttest	78.53	8.49	Follow-up
Externalizing Symptoms	Emotional Disclosure Drawing	Pretest	86.86	15.41	Posttest	54.40	5.61	Follow-up
	Floortime Play Therapy	Pretest	90.26	12.76	Posttest	52.60	5.43	Follow-up
	Control	Pretest	92.26	15.50	Posttest	70.46	7.84	Follow-up

As shown in Table 1, both intervention groups experienced marked improvements in emotion regulation and behavioral symptoms compared with the control group. In the emotional disclosure through drawing

group, the mean score for emotion regulation difficulties dropped sharply from 116.47 at pretest to 70.13 at posttest and remained low at follow-up (71.06). Internalizing and externalizing symptoms also declined notably, with internalizing scores falling from 88.06 to 50.93 and externalizing scores from 86.86 to 53.20 by follow-up. Similarly, in the Floortime play therapy group, emotion regulation difficulties decreased from 115.13 to 73.06 at posttest and stayed stable (74.20), while internalizing symptoms declined from 83.00 to 52.46 and externalizing symptoms from 90.26 to 52.60. In contrast, the control group showed only modest reductions, with emotion regulation difficulties decreasing slightly from 110.80 to 89.30 and internalizing and externalizing symptoms remaining relatively high at follow-up. These descriptive trends indicate that both interventions were effective in improving emotion regulation and reducing anxiety-related behavioral difficulties, with effects persisting over time.

Before conducting the main statistical analyses, all necessary assumptions for analysis of covariance (ANCOVA) were examined and confirmed. The normality of the data distribution was assessed using skewness, kurtosis, and the Shapiro–Wilk test, indicating that scores for all key variables were normally distributed. Homogeneity of variances across groups was verified with Levene’s test, which showed non-significant results, confirming equal error variances. The assumption of homogeneity of regression slopes was also met, as no significant interaction was found between the covariate (pretest scores) and the group variable. Additionally, linear relationships between the covariate and dependent variables were observed, and no outliers or influential data points were detected that could bias the results. These checks validated the suitability of the data for ANCOVA and ensured the reliability of the subsequent inferential analyses.

Table 2. Analysis of covariance (ANCOVA) results for emotion regulation difficulties, and multivariate ANCOVA for internalizing and externalizing symptoms

Effect	Variable	Sum of Squares	df	Mean Square	F	Sig.	Partial Eta Squared
Emotion Regulation Difficulties							
Corrected Model		12053.926	3	4017.975	11.334	.000	.453
Intercept		10215.139	1	10215.139	28.815	.000	.413
Pretest		304.192	1	304.192	0.858	.363	.020
Group		10582.907	2	5291.454	14.926	.001	.421
Error		14534.874	41	354.509			
Total		719253	45				
Internalizing Symptoms							
Corrected Model	Post-Internalizing	2460.862	4	615.216	15.219	.000	.603
Intercept	Post-Internalizing	2297.483	1	2297.483	56.836	.000	.587
Pre-Internalizing	Post-Internalizing	11.447	1	11.447	0.283	.598	.007
Pre-Externalizing	Post-Internalizing	29.032	1	29.032	0.718	.402	.018
Group	Post-Internalizing	2454.698	2	1227.349	30.363	.000	.603
Error	Post-Internalizing	1616.915	40	40.423			
Externalizing Symptoms							
Corrected Model	Post-Externalizing	2985.747	4	746.437	18.421	.000	.648
Intercept	Post-Externalizing	2579.153	1	2579.153	63.650	.000	.614
Pre-Internalizing	Post-Externalizing	91.125	1	91.125	2.249	.142	.053
Pre-Externalizing	Post-Externalizing	2.745	1	2.745	0.068	.796	.002
Group	Post-Externalizing	2960.518	2	1480.259	36.531	.000	.646
Error	Post-Externalizing	1620.831	40	40.521			

Table 2 shows the results of the ANCOVA for emotion regulation difficulties and the multivariate ANCOVA for internalizing and externalizing symptoms after controlling for pretest scores. For emotion regulation difficulties, the group effect was significant ($F = 14.926$, $p = .001$, $\eta^2 = .421$), indicating that the type of intervention contributed substantially to changes in regulation abilities beyond pretest differences. Similarly, multivariate analysis revealed significant group effects for both internalizing symptoms ($F =$

30.363, $p < .001$, $\eta^2 = .603$) and externalizing symptoms ($F = 36.531$, $p < .001$, $\eta^2 = .646$), demonstrating large effect sizes and meaningful reductions in behavioral problems across the intervention groups compared to the control. Pretest scores did not significantly predict posttest outcomes, confirming the effectiveness of the interventions independent of initial symptom severity. These findings support the robust impact of both emotional disclosure through drawing and family-based Floortime play therapy in improving emotional and behavioral functioning in children with anxiety symptoms.

Table 3. Bonferroni post hoc comparisons for emotion regulation difficulties and internalizing and externalizing symptoms

Dependent Variable	Stage	Group (I)	Group (J)	Std. Error	Mean Difference (I–J)	Sig.
Emotion Regulation Difficulties	Posttest	Emotional Disclosure Drawing	Floortime Play Therapy	2.5952	-2.9	.794
		Emotional Disclosure Drawing	Control	2.5952	-19.9	.000
		Floortime Play Therapy	Control	2.5952	-17.0	.000
	Follow-up	Emotional Disclosure Drawing	Floortime Play Therapy	2.9604	-3.13	.888
		Emotional Disclosure Drawing	Control	2.9604	-18.26	.000
		Floortime Play Therapy	Control	2.9604	-15.13	.000
Internalizing Symptoms	Posttest	Emotional Disclosure Drawing	Floortime Play Therapy	2.2952	-1.86	.100
		Emotional Disclosure Drawing	Control	2.2952	-14.5	.000
		Floortime Play Therapy	Control	2.2952	-12.4	.000
	Follow-up	Emotional Disclosure Drawing	Floortime Play Therapy	2.1839	-1.5	.100
		Emotional Disclosure Drawing	Control	2.1839	-16.2	.000
		Floortime Play Therapy	Control	2.1839	-14.6	.000
Externalizing Symptoms	Posttest	Emotional Disclosure Drawing	Floortime Play Therapy	2.3344	1.73	.100
		Emotional Disclosure Drawing	Control	2.3344	-16.06	.000
		Floortime Play Therapy	Control	2.3344	-15.8	.000
	Follow-up	Emotional Disclosure Drawing	Floortime Play Therapy	1.8295	0.6	.100
		Emotional Disclosure Drawing	Control	1.8295	-11.0	.000
		Floortime Play Therapy	Control	1.8295	-11.0	.000

Table 3 presents the Bonferroni post hoc comparisons among the three study groups for emotion regulation difficulties and internalizing and externalizing symptoms at posttest and follow-up. For emotion regulation difficulties, both the emotional disclosure and Floortime groups showed significantly lower posttest and follow-up scores compared with the control group ($p < .001$), while the difference between the two intervention groups was not significant at either time point ($p > .05$). Similarly, for internalizing symptoms, both active interventions produced significantly greater reductions compared with the control group at posttest and follow-up ($p < .001$), but no significant difference was observed between emotional disclosure and Floortime ($p > .05$). Externalizing symptoms followed the same pattern: both treatment groups improved significantly compared to the control group ($p < .001$), and no significant difference emerged between the two experimental interventions ($p > .05$). These findings confirm that while both interventions were highly effective relative to no treatment, their outcomes were comparable to each other in reducing emotion dysregulation and anxiety-related behavioral problems.

Discussion and Conclusion

The findings of this study demonstrated that both emotional disclosure through drawing and family-based Floortime play therapy produced significant improvements in children with anxiety symptoms, specifically by reducing emotion regulation difficulties and alleviating internalizing and externalizing behavioral problems. After controlling for pretest differences, both interventions outperformed the control group at posttest and follow-up, indicating robust and sustained therapeutic effects. Importantly, although each intervention significantly improved outcomes, there were no meaningful differences between the two active treatment groups, suggesting that both approaches are comparably effective in addressing the core emotional and behavioral difficulties of anxious children.

The strong reduction in emotion regulation difficulties aligns with the theoretical and empirical rationale underlying both interventions. Emotional disclosure through drawing provides children with a symbolic, developmentally appropriate medium to express distressing memories and unresolved affect, thereby reducing emotional avoidance and increasing self-awareness (16, 18). Visual representation of inner experiences helps reorganize negative emotions into coherent narratives, enhancing clarity and acceptance (17, 19). Similarly, the Floortime approach facilitates emotion regulation by fostering safe parent–child interactions, helping caregivers respond to affective cues and co-regulate emotional arousal (9, 11). Our findings that emotion regulation improved substantially in both groups echo previous studies showing that play-based, relationship-focused interventions strengthen self-regulation capacities in children with developmental and emotional difficulties (10, 13).

Improvements in internalizing symptoms—including anxiety, withdrawal, and somatic complaints—further support the utility of experiential therapies for anxious children. Children who engaged in expressive art were able to externalize fears and sadness, reducing internal tension and avoidance (17, 18). Past research shows that structured emotional disclosure can decrease symptoms of anxiety and depression by increasing tolerance for negative affect and promoting meaning-making (16). Likewise, the Floortime model’s emphasis on attunement and strengthening secure parent–child bonds likely helped reduce fearfulness and promote emotional safety, which are crucial for anxious children (9, 12). These results also align with the transdiagnostic perspective, which suggests that interventions targeting underlying affective mechanisms can reduce multiple domains of psychopathology (1, 4).

The observed decrease in externalizing symptoms is noteworthy because anxious children are often assumed to present primarily with internalizing difficulties; however, poor emotion regulation can also manifest as aggression and defiance when children are overwhelmed (2, 3). Our findings parallel studies showing that interventions enhancing emotional expression and parent–child relational quality can reduce oppositional behaviors and impulsivity (13, 20). The active involvement of parents in the Floortime program likely helped reduce maladaptive cycles of coercion and frustration, while art-based disclosure provided a non-threatening outlet for anger and irritability (9, 18).

Another important aspect is the sustained effectiveness observed at follow-up. Many interventions show initial improvements but fail to maintain gains once sessions end. The continued progress in both groups indicates that the skills and emotional insights acquired during therapy may have generalized to daily life. Floortime’s training of parents as emotional co-regulators could have contributed to ongoing support at

home (9, 15). Similarly, emotional disclosure through drawing may have enhanced children's internal coping strategies and ability to self-soothe when faced with distress (17, 19).

The comparable effectiveness of both methods is an important contribution to the literature. While Floortime is well-established in the treatment of autism and developmental disorders (11, 12), its use for anxiety and emotion dysregulation has been less systematically explored (13). Likewise, while art-based emotional disclosure has shown promise for aggression reduction and stress relief (15, 18), comparative trials with relational therapies have been scarce. Our findings support the adaptability of these modalities to anxiety-related emotional and behavioral issues and underscore the potential for non-pharmacological, developmentally attuned interventions to promote mental health in young populations (4, 5).

Furthermore, the family-based nature of Floortime appears to address relational vulnerabilities that can sustain anxiety. Studies have shown that high parental stress, unresolved trauma, and maladaptive emotional climates can intensify children's regulatory problems (7, 8). By coaching parents to tune into their child's emotional signals and maintain attuned interactions, Floortime interrupts these negative cycles. This aligns with earlier Iranian and international research demonstrating that parent-mediated interventions are crucial for sustainable child outcomes (9, 14).

In addition, the accessibility and cultural appropriateness of both interventions are notable. Drawing and play are natural communication tools for children across cultures and require minimal language complexity, making them highly adaptable in contexts where stigma or limited access to specialized services might hinder traditional therapy (17, 19). Prior research in Iranian populations has confirmed that both Floortime and art-based interventions can be effectively localized while preserving their therapeutic essence (9, 20). This cultural adaptability is essential for expanding child mental health care in diverse educational and clinical settings.

Despite the promising results, some limitations should be acknowledged. First, the study's sample size, while statistically adequate, was relatively small and drawn from a single geographic area, potentially limiting the generalizability of the findings. Children were selected using purposive sampling, which may introduce selection bias and reduce external validity. Second, the reliance on parent-report measures for internalizing and externalizing symptoms, while practical and widely used, might be influenced by parental perceptions and expectations, especially since parents were involved in one of the interventions. Including teacher reports or direct behavioral observations would strengthen future studies. Third, the follow-up period, although sufficient to observe short-term maintenance, was limited to a relatively brief post-intervention window; long-term durability of treatment effects remains unknown. Finally, while both interventions were manualized and structured, therapist effects and fidelity to the protocol were not independently assessed, leaving room for variability in implementation.

Future studies could build upon these findings by employing larger, more diverse samples drawn from multiple regions to improve generalizability and cultural representativeness. Longitudinal research with extended follow-up periods would be valuable to determine the persistence of treatment gains and to identify factors that predict long-term success. Incorporating multi-informant and multi-method assessments, including teacher ratings, direct observations, and physiological markers of emotion regulation, would add depth and reduce rater bias. Comparative studies could also explore hybrid models, integrating elements of Floortime's relational coaching with art-based emotional disclosure to examine potential synergistic effects.

Additionally, evaluating the cost-effectiveness and scalability of these approaches in school settings and community clinics would provide practical insights for broad implementation.

Practitioners working with anxious children may consider adopting both emotional disclosure through drawing and family-based Floortime as effective, non-invasive, and culturally flexible interventions. Floortime can empower caregivers by teaching them to become sensitive co-regulators and strengthen family emotional climates, while drawing-based disclosure can be easily implemented in individual or group settings to facilitate safe expression and processing of distress. Schools and mental health centers can integrate these interventions into preventive programs, offering support to children with early signs of anxiety before symptoms escalate. Training clinicians, educators, and parents in these approaches may enhance their capacity to nurture emotion regulation skills and reduce behavioral problems, ultimately promoting resilience and well-being in children.

Acknowledgments

The authors express their deep gratitude to all participants who contributed to this study.

Authors' Contributions

All authors equally contributed to this study.

Declaration of Interest

The authors of this article declared no conflict of interest.

Ethical Considerations

The study protocol adhered to the principles outlined in the Helsinki Declaration, which provides guidelines for ethical research involving human participants. This research was approved and registered by the Ethics Committee of the Islamic Azad University, Birjand Branch, under the ethical approval code IR.IAU.BIRJAND.REC.1403.028.

Transparency of Data

In accordance with the principles of transparency and open research, we declare that all data and materials used in this study are available upon request.

Funding

This research was carried out independently with personal funding and without the financial support of any governmental or private institution or organization.

References

1. Zitzmann J, Rombold-George L, Rosenbach C, Renneberg B. Emotion Regulation, Parenting, and Psychopathology: A Systematic Review. *Clinical Child and Family Psychology Review*. 2024;27(1):1-22. doi: 10.1007/s10567-023-00452-5.
2. Rajabi G, Abbasi Q. Relationship between self-criticism, social anxiety, and fear of failure with internalized shame in students. *Psychol Clin Consult Res*. 2011;1(2):171-82.

3. Kwon K, Hanrahan AR, Kupzyk K. Emotional Expressivity and Emotion Regulation: Relation to Academic Functioning Among Elementary School Children. *School Psychology Quarterly*. 2017;32(1):75-88. doi: 10.1037/spq0000166.
4. Pouladi S, Hasanshahi MM, Rabiei M, Bagheri N. The Effect of Unified Transdiagnostic Treatment on the Improvement of Internalizing Behavioral Problems, Emotional Regulation, and Empathy in Children with Anxiety Disorders. *Research in Cognitive and Behavioral Sciences*. 2022;12(1):141-64. doi: 10.22108/cbs.2022.133945.1655.
5. Masipour O-B. Effectiveness of Stress Inoculation Training on Emotional Regulation, Childbirth Self-Efficacy and Natural Childbirth Anxiety. *Aftj*. 2025;6(2):23-30. doi: 10.61838/kman.aftj.6.2.3.
6. Dragomir AM. The Dynamics of Child Anxiety as an Expression of Their Emotional Security in the Coparental System. *Univers Pedagogic*. 2024;83(3):85-90. doi: 10.52387/1811-5470.2024.3.13.
7. Bae S, Choi H-j. A Sequential Mediating Effects of Perfectionism, Suppression of Emotional Expression, and Parenting Stress in the Relationship Between Parentification and Housework Burnout in Women With Children. *Korean Association for Learner-Centered Curriculum and Instruction*. 2023;23(13):797-809. doi: 10.22251/jlcci.2023.23.13.797.
8. Hadiyan N, Mottaghi S, Rezapour-Mirsaleh Y. Investigating the Mediating Role of Maladaptive Schemas and Emotional Expression in the Relationship Between Childhood Trauma and Sexual Intimacy in Women From Low-Income Families. *Aftj*. 2023;4(5):610-32. doi: 10.61838/kman.aftj.4.5.35.
9. Abazari K, Malekpour M, Ghamarani A, Abedi A, Faramarzi S. Impact of Individual Differences Intervention (Floortime) Based on Parents' Expressed Emotion on Children's Social Skills With High-Functioning Autism Disorder. *Iranian Journal of Psychiatry and Clinical Psychology*. 2017;23(3):260-77. doi: 10.29252/nirp.ijpcp.23.3.260.
10. Asmeri Noubari F, Kafash Khormizi E, Tavakoli Dehghi Z, Silavi A, editors. The impact of Floortime play therapy (family-based) on enhancing children's social skills. *Second Conference on Psychology, Educational Sciences, Social Sciences, and Counseling*; 2023; Italy.
11. Şengül-Erdem H. DIR/Floortime: Otizm Spektrum Bozukluğunda Bir Erken Müdahale Modeli. *Kalem Uluslararası Eğitim Ve İnsan Bilimleri Dergisi*. 2021;11(1/20):21-39. doi: 10.23863/kalem.2021.177.
12. Bayat MR. Comparison of the Effectiveness of Applied Behavior Analysis Training, Floortime-Based Play Therapy, and Their Combination in Improving Communication and Social Skills of Children with Autism Spectrum Disorders in Andimeshk: *Doctoral Dissertation, Faculty of Educational Sciences and Psychology, Shahid Chamran University of Ahvaz*; 2016.
13. Roghani F, Jadidi M, Peymani J. The Effectiveness of Floortime Play Therapy on Improving Executive Functions and Cognitive Emotion Regulation in Children with Attention Deficit / Hyperactivity Disorder (ADHD). *International Journal of Education and Cognitive Sciences*. 2022;2(4):30-44. doi: 10.22034/injoeas.2022.160686.
14. Kohansal L, editor The impact of Floortime play therapy (family-based) on enhancing children's social skills. *Second Conference on Psychology, Educational Sciences, Social Sciences, and Counseling*; 2021; Italy.
15. Ebadi M, Mardani-Hamoooleh M, Seyedfatemi N, Ghaljeh M, Haghani SH. Effects of Written Emotional Disclosure on the Stress of Mothers of Children With Autism. *Iran Journal of Nursing*. 2021;34(131):19-30. doi: 10.52547/ijn.34.131.19.
16. Bottoms BL, Peter-Hagene LC, Epstein MA, Wiley T, Reynolds CE, Rudnicki AG. Abuse Characteristics and Individual Differences Related to Disclosing Childhood Sexual, Physical, and Emotional Abuse and Witnessed Domestic Violence. *Journal of Interpersonal Violence*. 2014;31(7):1308-39. doi: 10.1177/0886260514564155.
17. Liu Z, Yao C, Ying F. Effects of Digital Art Emotional Expression Therapy on the Rehabilitation of Children With Autism Spectrum Disorder. 2024. doi: 10.54941/ahfe1004556.
18. Anjam SZ, Dortaj F. Investigating the effectiveness of emotional disclosure in the form of painting on reducing aggression in female children. *Journal of Advanced Pharmacy Education and Research*. 2020;10(4):120-6.
19. Safikhani F. The effectiveness of grammatical mental imagery with cognitive processing on self-efficacy, emotional processing and spirituality in mothers of students with autism spectrum disorder. *International Journal of Education and Cognitive Sciences*. 2022;3(2):12-22. doi: 10.22034/injoeas.2022.160609.

20. Mohammadi Keshka S, Abbas SM, Abolghasemi MK. The effect of combining group play therapy with painting on internalizing and externalizing problems in children with separation anxiety disorder. *Journal of Exceptional Education and Training*. 2022;5(171):24-34.